arXiv:1404.0886v2 [math.CV] 4 Apr 2014

$(\alpha, \beta, \lambda, \delta, m, \Omega)_p-$NEIGHBORHOOD FOR SOME FAMILIES OF ANALYTIC AND MULTIVALENT FUNCTIONS

HALIT ORHAN

Abstract. In the present investigation, we give some interesting results related with neighborhoods of $p-$valent functions. Relevant connections with some other recent works are also pointed out.

1. INTRODUCTION AND DEFINITIONS

Let A demonstrate the family of functions $f(z)$ of the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n$$

which are analytic in the open unit disk $\mathcal{U} = \{z \in \mathbb{C} : |z| < 1\}$.

We denote by $\mathcal{A}_p(n)$ the class of functions $f(z)$ normalized by

$$f(z) = z^p + \sum_{k=n}^{\infty} a_{k+p} z^{k+p} \quad (n, p \in \mathbb{N} := \{1, 2, 3, ...\}) \tag{1.1}$$

which are analytic and p-valent in $\mathcal{U}$.

Upon differentiating both sides of (1.1) m times with respect to z, we have

$$f^{(m)}(z) = \frac{p!}{(p-m)!} z^{p-m} + \sum_{k=n}^{\infty} \frac{(k+p)!}{(k+p-m)!} a_{k+p} z^{k+p-m} \tag{1.2}$$

$$(n, p \in \mathbb{N}; m \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}; p > m).$$

We show by $\mathcal{A}_p(n, m)$ the class of functions of the form (1.2) which are analytic and p-valent in $\mathcal{U}$.

1991 *Mathematics Subject Classification.* 30C45.

Key words and phrases. $p-$valent functions, inclusion relations, neighborhood properties, Salagean differential operator, Miller and Mocanu's lemma.

The concept of neighborhood for $f(z) \in \mathcal{A}$ was first given by Goodman [7]. The concept of δ-neighborhoods $N_\delta(f)$ of analytic functions $f(z) \in \mathcal{A}$ was first studied by Ruscheweyh [8]. Walker [12], defined a neighborhood of analytic functions having positive real part. Later, Owa *et al.*[13] generalized the results given by Walker. In 1996, Altıntaş and Owa [14] gave (n, δ)-neighborhoods for functions $f(z) \in \mathcal{A}$ with negative coefficients. In 2007, (n, δ)-neighborhoods for p-valent functions with negative coefficients were considered by Srivastava *et al.* [4], and Orhan [5]. Very recently, Orhan *et al.*[1], introduced a new definition of (n, δ)-neighborhood for analytic functions $f(z) \in \mathcal{A}$. Orhan *et al.*'s [1] results were generalized for the functions $f(z) \in \mathcal{A}$ and $f(z) \in \mathcal{A}_p(n)$ by many author (see, [6, 9, 10, 15]).

In this paper, we introduce the neighborhoods $(\alpha, \beta, \lambda, m, \delta, \Omega)_p - N(g)$ and $(\alpha, \beta, \lambda, m, \delta, \Omega)_p - M(g)$ of a function $f^{(m)}(z)$ when $f(z) \in \mathcal{A}_p(n)$.

Using the Salagean derivative operator [3]; we can write the following equalities for the function $f^{(m)}(z)$ given by

$$D^0 f^{(m)}(z) = f^{(m)}(z)$$

$$D^1 f^{(m)}(z) = \frac{z}{(p-m)}\left(f^{(m)}(z)\right)' = \frac{p!}{(p-m)!}z^{p-m} + \sum_{k=n}^{\infty} \frac{(k+p-m)(k+p)!}{(p-m)(k+p-m)!}a_{k+p}z^{k+p-m}$$

$$D^2 f^{(m)}(z) = D(Df^{(m)}(z)) = \frac{p!}{(p-m)!}z^{p-m} + \sum_{k=n}^{\infty} \frac{(k+p-m)^2(k+p)!}{(p-m)^2(k+p-m)!}a_{k+p}z^{k+p-m}$$

.....

$$D^\Omega f^{(m)}(z) = D(D^{\Omega-1} f^{(m)}(z)) = \frac{p!}{(p-m)!}z^{p-m} + \sum_{k=n}^{\infty} \frac{(k+p-m)^\Omega(k+p)!}{(p-m)^\Omega(k+p-m)!}a_{k+p}z^{k+p-m}.$$

We define $F : \mathcal{A}_p(n, m) \to \mathcal{A}_p(n, m)$ such that

$$F(f^{(m)}(z)) = (1-\lambda)\left(D^\Omega f^{(m)}(z)\right) + \frac{\lambda z}{(p-m)}\left(D^\Omega f^{(m)}(z)\right)'$$

$$= \frac{p!}{(p-m)!}z^{p-m} + \sum_{k=n}^{\infty} \frac{(k+p)!(k+p-m)^\Omega(1+\lambda k(p-m)^{-1})}{(p-m)^\Omega(k+p-m)!}a_{k+p}z^{k+p-m} \tag{1.3}$$

$$(0 \leq \lambda \leq 1;\ \Omega, m \in \mathbb{N}_0;\ p > m).$$

Let $F(\lambda, m, \Omega)$ denote class of functions of the form (1.3) which are analytic in $\mathcal{U}$.

For $f, g \in F(\lambda, m, \Omega)$, f said to be $(\alpha, \beta, \lambda, m, \delta, \Omega)_p$−neighborhood for g if it satisfies

$$\left| \frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} \right| < \delta \quad (z \in \mathcal{U})$$

for some $-\pi \leq \alpha - \beta \leq \pi$ *and* $\delta > \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$. We show this neighborhood by $(\alpha, \beta, \lambda, m, \delta, \Omega)_p - N(g)$.

Also, we say that $f \in (\alpha, \beta, \lambda, m, \delta, \Omega)_p - M(g)$ if it satisfies

$$\left| \frac{e^{i\alpha} F(f^{(m)}(z))}{z^{p-m}} - \frac{e^{i\beta} F(g^{(m)}(z))}{z^{p-m}} \right| < \delta \quad (z \in \mathcal{U})$$

for some $-\pi \leq \alpha - \beta \leq \pi$ *and* $\delta > \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$.

We give some results for functions belonging to $(\alpha, \beta, \lambda, m, \delta, \Omega)_p - N(g)$ and $(\alpha, \beta, \lambda, m, \delta, \Omega)_p - M(g)$.

2. Main Results

Now we can establish our main results.

Theorem 2.1. *If* $f \in F(\lambda, m, \Omega)$ *satisfies*

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!} \left| e^{i\alpha} a_{k+p} - e^{i\beta} b_{k+p} \right|$$

$$\leq \delta - \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]} \tag{2.1}$$

for some $-\pi \leq \alpha - \beta \leq \pi$; $p > m$ *and* $\delta > \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$, *then* $f \in (\alpha, \beta, \lambda, m, \delta, \Omega)_p - N(g)$.

Proof. By virtue of (1.3), we can write

$$\left| \frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} \right| =$$

$$\left|\frac{p!(p-m)}{(p-m)!}e^{i\alpha}+e^{i\alpha}\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}a_{k+p}z^{k}-\frac{p!(p-m)}{(p-m)!}e^{i\beta}\right.$$
$$\left.-e^{i\beta}\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}b_{k+p}z^{k}\right|$$

$$< \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$$
$$+\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p}\right|.$$

If

$$\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p}\right|$$
$$\leq \delta-\frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]},$$

then we observe that

$$\left|\frac{e^{i\alpha}F'(f^{(m)}(z))}{z^{p-m-1}}-\frac{e^{i\beta}F'(g^{(m)}(z))}{z^{p-m-1}}\right|<\delta \quad (z\in\mathcal{U}).$$

Thus, $f\in(\alpha,\beta,\lambda,m,\delta,\Omega)_p-N(g)$. This evidently completes the proof of Theorem 2.1. □

Remark 2.2. *In its special case when*

$$m=\Omega=\lambda=\alpha=0 \text{ and } \beta=\alpha, \tag{2.2}$$

in Theorem 2.1 yields a result given earlier by Altuntaş et al. ([9] *p.3, Theorem 1).*

We give an example for Theorem 2.1.

Example 2.1. *For given*

$$g(z)=\frac{p!}{(p-m)!}z^{p-m}+\sum_{k=n}^{\infty}B_{k+p}(\alpha,\beta,\lambda,m,\delta,\Omega)z^{k+p-m}\in F(\lambda,m,\Omega)$$
$$(n,p\in\mathbb{N}=\{1,2,3,...\};\ p>m;\ \Omega,m\in\mathbb{N}_0)$$

we consider

$$f(z)=\frac{p!}{(p-m)!}z^{p-m}+\sum_{k=n}^{\infty}A_{k+p}(\alpha,\beta,\lambda,m,\delta,\Omega)z^{k+p-m}\in F(\lambda,m,\Omega)$$
$$(n,p\in\mathbb{N}=\{1,2,3,...\};\ p>m;\ \Omega,m\in\mathbb{N}_0)$$

with

$$A_{k+p}=\frac{(p-m)^{\Omega}\{\delta-\frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}\}(k+p-m)!(n+p-1)}{(1+\lambda k(p-m)^{-1})(k+p-m)^{\Omega+1}(k+p-1)!(k+p)^{2}(k+p-1)}e^{-i\alpha}+e^{i(\beta-\alpha)}B_{k+p}.$$

Then we have that

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!} \left|e^{i\alpha}A_{k+p} - e^{i\beta}B_{k+p}\right| =$$

$$(n+p-1)\left(\delta - \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}\right)\sum_{k=n}^{\infty}\frac{1}{(k+p-1)(k+p)}. \tag{2.3}$$

Finally, in view of the telescopic series, we obtain

$$\begin{aligned}\sum_{k=n}^{\infty}\frac{1}{(k+p-1)(k+p)} &= \lim_{\varsigma\longrightarrow\infty}\sum_{k=n}^{\varsigma}\left[\frac{1}{k+p-1}-\frac{1}{k+p}\right] \\ &= \lim_{\varsigma\longrightarrow\infty}\left[\frac{1}{n+p-1}-\frac{1}{\zeta+p}\right] \\ &= \frac{1}{n+p-1}.\end{aligned} \tag{2.4}$$

Using (2.4) in (2.3), we get

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!} \left|e^{i\alpha}A_{k+p} - e^{i\beta}B_{k+p}\right|$$

$$= \delta - \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}.$$

Therefore, we say that $f \in (\alpha,\beta,\lambda,m,\delta,\Omega)_p - N(g)$.

Also, Theorem 1 gives us the following corollary.

Corollary 2.3. *If $f \in F(\lambda,m,\Omega)$ satisfies*

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!} \left||a_{k+p}| - |b_{k+p}|\right|$$

$$\leq \delta - \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$$

for some $-\pi \leq \alpha-\beta \leq \pi$ and $\delta > \frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}$, and $\arg(a_{k+p}) - \arg(b_{k+p}) = \beta-\alpha$ $(n,p \in \mathbb{N} = \{1,2,3,...\};\ m \in \mathbb{N}_0,\ p > m)$, then $f \in (\alpha,\beta,\lambda,m,\delta,\Omega)_p - N(g)$.

Proof. By theorem (2.1), we see the inequality (2.1) which implies that $f \in (\alpha,\beta,\lambda,m,\delta,\Omega)_p - N(g)$.

Since $\arg(a_{k+p}) - \arg(b_{k+p}) = \beta-\alpha$, if $\arg(a_{k+p}) = \alpha_{k+p}$, we see $\arg(b_{k+p}) = \alpha_{k+p} - \beta + \alpha$. Therefore,

$$e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p} = e^{i\alpha}\left|a_{k+p}\right|e^{i\alpha_{k+p}} - e^{i\beta}\left|b_{k+p}\right|e^{i(\alpha_{k+p}-\beta+\alpha)} = (|a_{k+p}| - |b_{k+p}|)e^{i(\alpha_{k+p}+\alpha)}$$

implies that

$$\left|e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p}\right|=\left||a_{k+p}|-|b_{k+p}|\right|. \tag{2.5}$$

Using (2.5) in (2.1) the proof of the corollary is complete.

Next, we can prove the following theorem. □

Theorem 2.4. *If $f\in F(\lambda,m,\Omega)$ satisfies*

$$\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p}\right|\leq\delta-\frac{p!}{(p-m)!}\sqrt{2[1-\cos(\alpha-\beta)]}\quad(z\in\mathcal{U}).$$

for some $-\pi\leq\alpha-\beta\leq\pi; p>m$ and $\delta>\frac{p!}{(p-m)!}\sqrt{2[1-\cos(\alpha-\beta)]}$ then $f\in(\alpha,\beta,\lambda,m,\delta,\Omega)_p-M(g)$.

The proof of this teorem is similar with Theorem 2.1.

Corollary 2.5. *If $f\in F(\lambda,m,\Omega)$ satisfies*

$$\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left||a_{k+p}|-|b_{k+p}|\right|\leq\delta-\frac{p!}{(p-m)!}\sqrt{2[1-\cos(\alpha-\beta)]}\quad(z\in\mathcal{U}).$$

for some $-\pi\leq\alpha-\beta\leq\pi$; $p>m$ and $\delta>\frac{p!}{(p-m)!}\sqrt{2[1-\cos(\alpha-\beta)]}$ and $\arg(a_{k+p})-\arg(b_{k+p})=\beta-\alpha$, then $f\in(\alpha,\beta,\lambda,m,\delta,\Omega)_p-M(g)$.

Our next result as follows.

Theorem 2.6. *If $f\in(\alpha,\beta,\lambda,m,\delta,\Omega)_p-N(g), 0\leq\alpha<\beta\leq\pi$; $p>m$ and $\arg(e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p})=k\phi$, then*

$$\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p}\right|\leq\delta-\frac{p!}{(p-m-1)!}(\cos\alpha-\cos\beta).$$

Proof. For $f\in(\alpha,\beta,\lambda,m,\delta,\Omega)_p-N(g)$, we have

$$\begin{aligned}&\left|\frac{e^{i\alpha}F'(f^{(m)}(z))}{z^{p-m-1}}-\frac{e^{i\beta}F'(g^{(m)}(z))}{z^{p-m-1}}\right|\\ =&\left|\frac{p!(e^{i\alpha}-e^{i\beta})}{(p-m-1)!}+\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}(e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p})z^k\right|\end{aligned}$$

$$=\left|\frac{p!(e^{i\alpha}-e^{i\beta})}{(p-m-1)!}+\sum_{k=n}^{\infty}\frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}(e^{i\alpha}a_{k+p}-e^{i\beta}b_{k+p})e^{ik\phi}z^k\right|<\delta.$$

Let us consider z such that $\arg z = -\phi$. Then $z^k = |z|^k e^{-ik\phi}$. For such a point $z \in \mathcal{U}$, we see that

$$\begin{aligned}
&\left|\frac{e^{i\alpha}F'(f(z))}{z^{p-m-1}} - \frac{e^{i\beta}F'(g(z))}{z^{p-m-1}}\right| \\
&= \left|\frac{p!(e^{i\alpha}-e^{i\beta})}{(p-m-1)!} + \sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right||z|^k\right| \\
&= \left[\left(\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right||z|^k + \frac{p!(\cos\alpha-\cos\beta)}{(p-m-1)!}\right)^2\right. \\
&\qquad \left. + \left(\frac{p!(\sin\alpha-\sin\beta)}{(p-m-1)!}\right)^2\right]^{\frac{1}{2}} < \delta.
\end{aligned}$$

This implies that

$$\left(\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right||z|^k + \frac{p!(\cos\alpha-\cos\beta)}{(p-m-1)!}\right)^2 < \delta^2,$$

or

$$\frac{p!}{(p-m-1)!}(\cos\alpha-\cos\beta)+\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right||z|^k < \delta$$

for $z \in \mathcal{U}$. Letting $|z| \longrightarrow 1^-$, we have that

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega+1}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right| \leq \delta - \frac{p!}{(p-m-1)!}(\cos\alpha-\cos\beta).$$

□

Remark 2.7. *Applying the parametric substitutions listed in (2.2), Theorem 2.4 and 2.6 would yield a set of known results due to Altuntaş et al. ([9] p.5, Theorem 4; p.6, Theorem 7).*

Theorem 2.8. *If* $f \in (\alpha, \beta, \lambda, m, \delta, \Omega)_p - M(g), 0 \leq \alpha < \beta \leq \pi$ *and* $\arg(e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}) = k\phi$, *then*

$$\sum_{k=n}^{\infty} \frac{(k+p-m)^{\Omega}(k+p)!(1+\lambda k(p-m)^{-1})}{(p-m)^{\Omega}(k+p-m)!}\left|e^{i\alpha}a_{k+p} - e^{i\beta}b_{k+p}\right| \leq \delta + \frac{p!}{(p-m-1)!}(\cos\beta-\cos\alpha).$$

The proof of this theorem is similar with Theorem 2.6.

Remark 2.9. *Taking* $\lambda = \alpha = \Omega = m = 0$, $\beta = \alpha$ *and* $p = 1$,*in Theorem 2.8, we arrive at the following Theorem due to Orhan et al.*[1].

Theorem 2.10. *If* $f \in (\alpha, \delta) - N(g)$ *and* $\arg(a_n - e^{i\alpha}b_n) = (n-1)\varphi \quad (n = 2, 3, 4, ...)$, *then*

$$\sum_{n=2}^{\infty} n\left|a_n - e^{i\alpha}b_n\right| \leq \delta + \cos\alpha - 1.$$

We give an application of following lemma due to Miller and Mocanu [2] *(see also,* [11]*).*

Lemma 2.1. *Let the function*

$$w(z) = b_n z^n + b_{n+1} z^{n+1} + b_{n+2} z^{n+2} + ... \quad (n \in \mathcal{U})$$

be regular in $\mathcal{U}$ *with* $w(z) \neq 0$, $(n \in \mathcal{U})$. *If* $z_0 = r_0 e^{i\theta_0}$ $(r_0 < 1)$ *and* $|w(z_0)| = \max_{|z| \leq r_0} |w(z)|$, *then* $z_0 w'(z_0) = q w(z_0)$ *where* q *is real and* $q \geq n \geq 1$.

Applying the above lemma, we derive

Theorem 2.11. *If* $f \in F(\lambda, m, \Omega)$ *satisfies*

$$\left| \frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} \right| < \delta(p+n-m) - \frac{p!}{(p-m-1)!} \sqrt{2[1-\cos(\alpha-\beta)]}$$

for some $-\pi \leq \alpha - \beta \leq \pi$; $p > m$ *and* $\delta > \left(\frac{p!}{(p+n-m)(p-m-1)!} \right) \sqrt{2[1-\cos(\alpha-\beta)]}$, *then*

$$\left| \frac{e^{i\alpha} F(f^{(m)}(z))}{z^{p-m}} - \frac{e^{i\beta} F(g^{(m)}(z))}{z^{p-m}} \right| < \delta + \frac{p!}{(p-m)!} \sqrt{2[1-\cos(\alpha-\beta)]} \quad (z \in \mathcal{U}).$$

Proof. Let us define $w(z)$ by

$$\frac{e^{i\alpha} F(f^{(m)}(z))}{z^{p-m}} - \frac{e^{i\beta} F(g^{(m)}(z))}{z^{p-m}} = \frac{p!}{(p-m)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z). \tag{2.6}$$

Then $w(z)$ is analytic in $\mathcal{U}$ and $w(0) = 0$. By logarithmic differentiation, we get from (2.6) that

$$\frac{e^{i\alpha} F'(f^{(m)}(z)) - e^{i\beta} F'(g^{(m)}(z))}{e^{i\alpha} F(f^{(m)}(z)) - e^{i\beta} F(g^{(m)}(z))} - \frac{p-m}{z} = \frac{\delta w\prime(z)}{\frac{p!}{(p-m)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z)}.$$

Since

$$\frac{e^{i\alpha} F'(f^{(m)}(z)) - e^{i\beta} F'(g^{(m)}(z))}{z^{p-m} \left(\frac{p!}{(p-m)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z) \right)} = \frac{p-m}{z} + \frac{\delta w\prime(z)}{\frac{p!}{(p-m)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z)},$$

we see that

$$\frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} = \frac{p!}{(p-m-1)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z) \left(p - m + \frac{z w\prime(z)}{w(z)} \right).$$

This implies that

$$\left| \frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} \right| = \left| \frac{p!}{(p-m-1)!}(e^{i\alpha} - e^{i\beta}) + \delta w(z) \left(p - m + \frac{z w\prime(z)}{w(z)} \right) \right|.$$

We claim that

$$\left| \frac{e^{i\alpha} F'(f^{(m)}(z))}{z^{p-m-1}} - \frac{e^{i\beta} F'(g^{(m)}(z))}{z^{p-m-1}} \right| < \delta(p-m+n) - \frac{p!}{(p-m-1)!} \sqrt{2[1-\cos(\alpha-\beta)]}$$

in $\mathcal{U}$.

Otherwise, there exists a point $z_0 \in \mathcal{U}$ such that $z_0 w'(z_0) = q w(z_0)$ (by Miller and Mocanu's Lemma) where $w(z_0) = e^{i\theta}$ and $q \geq n \geq 1$.

Therefore, we obtain that

$$\begin{aligned}\left|\frac{e^{i\alpha}F'(f^{(m)}(z))}{z_0^{p-m-1}}-\frac{e^{i\beta}F'(g^{(m)}(z))}{z_0^{p-m-1}}\right| &= \left|\frac{p!}{(p-m-1)!}(e^{i\alpha}-e^{i\beta})+\delta e^{i\theta}\,(p-m+q)\right| \\ &\geq \delta\,(p+q-m)-\left|\frac{p!}{(p-m-1)!}(e^{i\alpha}-e^{i\beta})\right| \\ &\geq \delta\,(p+n-m)-\frac{p!}{(p-m-1)!}\sqrt{2[1-\cos(\alpha-\beta)]}.\end{aligned}$$

This contradicts our condition in Theorem 2.11.

Hence, there is no $z_0 \in \mathcal{U}$ such that $|w(z_0)| = 1$. This means that $|w(z)| < 1$ for all $z \in \mathcal{U}$.

Thus, have that

$$\begin{aligned}\left|\frac{e^{i\alpha}F(f^{(m)}(z))}{z^{p-m}}-\frac{e^{i\beta}F(g^{(m)}(z))}{z^{p-m}}\right| &= \left|\frac{p!}{(p-m)!}(e^{i\alpha}-e^{i\beta})+\delta w(z)\right| \\ &\leq \frac{p!}{(p-m)!}\left|e^{i\alpha}-e^{i\beta}\right|+\delta\,|w(z)| \\ &< \delta+\frac{p!}{(p-m)!}\sqrt{2[1-\cos(\alpha-\beta)]}.\end{aligned}$$

Upon setting $m = 0$, $\alpha = \varphi, \wp = F$ and $\beta = \alpha$ in Theorem 2.11, we have the following corollary given by Sağsöz et al.[6]. □

Corollary 2.12. If $f \in \wp(\Omega, \lambda)$ satisfies

$$\left|\frac{e^{i\alpha}\wp'(f(z))}{z^{p-1}}-\frac{e^{i\beta}\wp'(g(z))}{z^{p-1}}\right| < \delta(p+n)-p\sqrt{2[1-\cos(\varphi-\alpha)]}$$

for some $-\pi \leq \alpha-\beta \leq \pi$;*and* $\delta > \left(\frac{p}{(p+n)}\right)\sqrt{2[1-\cos(\alpha-\beta)]}$, then

$$\left|\frac{e^{i\alpha}\wp(f(z))}{z^{p}}-\frac{e^{i\beta}\wp(g(z))}{z^{p}}\right| < \delta+\sqrt{2[1-\cos(\varphi-\alpha)]} \quad (z \in \mathcal{U})\,.$$

□

References

[1] H. Orhan, E. Kadıoğlu and S. Owa, $(\alpha, \delta)-$Neighborhood for certain analytic functions, International Symposium on Geometric Function Theory and Applications (Edited by S. Owa and Y. Polatoğlu), T. C. Istanbul Kultur University Publ., 2008, 207-213.

[2] Miller, S.S and Mocanu, P.T, Second order differential inequalities in the complex plane, J. Math. Anal.Appl., 65, 289-305, (1978).

[3] G. S. Sălăgean, *Subclasses of univalent functions*, Complex analysis –Proc. 5th Rom.-Finn. Semin., Bucharest 1981, Part 1, Lect. Notes Math. 1013 (1983) 362-372.

[4] H. M. Srivastava and H. Orhan, Coefficient inequalities and inclusion relations for some families of analytic and multivalent functions, Appl. Math. Lett., 20 (2007), 686-691.

[5] H. Orhan, Neighborhoods of a certain class of p-valent functions with negative coefficients defined by using a differential operator, Math. Ineq. Appl. Vol. 12, Number 2, (2009), 335-349.

[6] F. Sağsöz, and M. Kamali, $(\varphi, \alpha, \delta, \lambda, \Omega)_p$-Neighborhood for some classes of multivalent functions, J. Ineq. Appl., (2013), 2013:152.

[7] Goodman, A.W, Univalent functions and nonanalytic curves, Proc. Amer. Math. Soc.,8, 598-601, (1957).

[8] Ruscheweyh, S., Neighborhoods of univalent functions,Proc. Amer. Math. Soc., 81, 521-527, (1981).

[9] F. Altuntaş, S. Owa and M. Kamali, $(\alpha, \delta)_p$-Neighborhood for certain class of multivalent functions, PanAmer. Math. J., Vol. 19, (2009), No. 235-46.

[10] Frasin, B. A., (α, β, δ)-Neighborhood for certain analytic functions with negative coefficients, Eur. J. Pure Appl.Math., 4(1), 14-19 (2011).

[11] I. S. Jack, Functions starlikenes and convex of order α, J. London Math. Soc. 2(3) (1971), 469-474.

[12] Walker, J. B., A note on neighborhoods of analytic functions having positive real part, Int. J. Math. Math. Sci., 13, 425–430, (1990).

[13] Owa, S., Saitoh, H. and Nunokawa, M., Neighborhoods of certain analytic functions, Appl. Math. Lett., 6, 73-77, (1993).

[14] Altınta̧s, O. and Owa, S., Neighborhood for certain analytic functions with negative coefficients, Int. J. Math. Math. Sci. 19, 797-800, (1996).

[15] Kugita, K., Kuroki, K. and Owa, S., (α, β)-Neighborhood for functions associated with Salagean differential operator and Alexander integral operator, Int. J. Math. Anal. Vol. 4, 2010, No. 5, 211-220.

Department of Mathematics, Faculty of Science, Ataturk University, 25240 Erzurum, Turkey.

E-mail address: orhanhalit607@gmail.com